# Recovery of algebraic-exponential data from moments

#### Jean B. Lasserre

LAAS-CNRS and Institute of Mathematics, Toulouse, France

#### TERRY FEST, LIMOGES, May 2015

#### \* Part of this work is joint with M. Putinar

A (1) > (1) > (1)

#### Motivation

#### Integrals

$$y \mapsto \int_{\{\mathbf{x}: g(x) \leq y\}} h(x) \, dx$$
 and  $g \mapsto \int_{\{\mathbf{x}: g(x) \leq 1\}} h(x) \, dx$ ,

with Positively Homogeneous Functions (PHF)

- Some properties (convexity, polarity)
- Sub-level sets of minimum volume containing  $\mathbf{K} \subset \mathbb{R}^n$
- Exact reconstruction from moments
- Recovery of the defining function of a semi-algebraic set

・ 同 ト ・ ヨ ト ・ ヨ ト

- Motivation
- Integrals

$$\mathbf{y} \mapsto \int_{\{\mathbf{x}: g(x) \leq \mathbf{y}\}} \mathbf{h}(x) \, dx \quad \text{and} \quad \mathbf{g} \mapsto \int_{\{\mathbf{x}: g(x) \leq 1\}} \mathbf{h}(x) \, dx,$$

- Some properties (convexity, polarity)
- Sub-level sets of minimum volume containing  $\mathbf{K} \subset \mathbb{R}^n$
- Exact reconstruction from moments
- Recovery of the defining function of a semi-algebraic set

・ 同 ト ・ ヨ ト ・ ヨ ト

- Motivation
- Integrals

$$\mathbf{y} \mapsto \int_{\{\mathbf{x}: \mathbf{g}(x) \leq \mathbf{y}\}} \mathbf{h}(x) \, dx \quad \text{and} \quad \mathbf{g} \mapsto \int_{\{\mathbf{x}: \mathbf{g}(x) \leq 1\}} \mathbf{h}(x) \, dx,$$

- Some properties (convexity, polarity)
- Sub-level sets of minimum volume containing  $\mathsf{K} \subset \mathbb{R}^n$
- Exact reconstruction from moments
- Recovery of the defining function of a semi-algebraic set

・ 回 ト ・ ヨ ト ・ ヨ ト

- Motivation
- Integrals

$$\mathbf{y} \mapsto \int_{\{\mathbf{x}: \mathbf{g}(x) \leq \mathbf{y}\}} \mathbf{h}(x) \, dx \quad \text{and} \quad \mathbf{g} \mapsto \int_{\{\mathbf{x}: \mathbf{g}(x) \leq 1\}} \mathbf{h}(x) \, dx,$$

- Some properties (convexity, polarity)
- Sub-level sets of minimum volume containing  $\mathbf{K} \subset \mathbb{R}^n$
- Exact reconstruction from moments
- Recovery of the defining function of a semi-algebraic set

(本間) (本語) (本語)

- Motivation
- Integrals

$$\mathbf{y} \mapsto \int_{\{\mathbf{x}: \mathbf{g}(x) \leq \mathbf{y}\}} \mathbf{h}(x) \, dx \quad \text{and} \quad \mathbf{g} \mapsto \int_{\{\mathbf{x}: \mathbf{g}(x) \leq 1\}} \mathbf{h}(x) \, dx,$$

- Some properties (convexity, polarity)
- Sub-level sets of minimum volume containing  $\mathbf{K} \subset \mathbb{R}^n$
- Exact reconstruction from moments
- Recovery of the defining function of a semi-algebraic set

・ 同 ト ・ ヨ ト ・ ヨ ト …

- Motivation
- Integrals

$$\mathbf{y} \mapsto \int_{\{\mathbf{x}: \mathbf{g}(x) \leq \mathbf{y}\}} \mathbf{h}(x) \, dx \quad \text{and} \quad \mathbf{g} \mapsto \int_{\{\mathbf{x}: \mathbf{g}(x) \leq 1\}} \mathbf{h}(x) \, dx,$$

- Some properties (convexity, polarity)
- Sub-level sets of minimum volume containing  $\mathbf{K} \subset \mathbb{R}^n$
- Exact reconstruction from moments
- Recovery of the defining function of a semi-algebraic set

・ 同 ト ・ ヨ ト ・ ヨ ト …

-

#### Reconstruction of a shape $\mathbf{K} \subset \mathbb{R}^n$ (convex or not)

from knowledge of finitely many moments

$$\mathbf{y}_{\alpha} = \int_{\mathbf{K}} x_1^{\alpha_1} \cdots x_n^{\alpha_n} \, dx, \qquad \alpha \in \mathbb{N}_d^n,$$

for some integer *d*, is a difficult and challenging problem!

#### EXACT recovery of **K**

from  $y = (y_{\alpha}), \alpha \in \mathbb{N}^{n}_{d}$ , is even more difficult and challenging!

イロト イポト イヨト イヨト 三日

#### Reconstruction of a shape $\mathbf{K} \subset \mathbb{R}^n$ (convex or not)

from knowledge of finitely many moments

$$\mathbf{y}_{\alpha} = \int_{\mathbf{K}} x_1^{\alpha_1} \cdots x_n^{\alpha_n} \, dx, \qquad \alpha \in \mathbb{N}_d^n,$$

for some integer *d*, is a difficult and challenging problem!

#### EXACT recovery of K

from  $\mathbf{y} = (\mathbf{y}_{\alpha}), \alpha \in \mathbb{N}^{n}_{\mathbf{d}}$ , is even more difficult and challenging!

ヘロト ヘアト ヘビト ヘビト

#### Examples of exact recovery:

- Quadrature (planar) Domains in (ℝ<sup>2</sup>) (Gustafsson, He, Milanfar and Putinar (Inverse Problems, 2000))
   via an exponential transform
- Convex Polytopes (in ℝ<sup>n</sup>) (Gravin, Lasserre, Pasechnik and Robins (Discrete & Comput. Geometry (2012))
   Use Brion-Barvinok-Khovanski-Lawrence-Pukhlikov moment formula for projections ∫<sub>P</sub> ⟨c, x⟩<sup>j</sup> dx combined with a Prony-type method to recover the vertices of *P*.
- and extension to Non convex polyhedra by Pasechnik et al.
  - via inversion of Fantappié transform

#### Examples of exact recovery:

- Quadrature (planar) Domains in (ℝ<sup>2</sup>) (Gustafsson, He, Milanfar and Putinar (Inverse Problems, 2000))
   via an exponential transform
- Convex Polytopes (in ℝ<sup>n</sup>) (Gravin, Lasserre, Pasechnik and Robins (Discrete & Comput. Geometry (2012))
   Use Brion-Barvinok-Khovanski-Lawrence-Pukhlikov moment formula for projections ∫<sub>P</sub> ⟨c, x⟩<sup>j</sup> dx combined with a Prony-type method to recover the vertices of *P*.
- and extension to Non convex polyhedra by Pasechnik et al.
  - via inversion of Fantappié transform

#### Approximate recovery can de done in multi-dimensions

(Cuyt, Golub, Milanfar and Verdonk, 2005) via :

- (multi-dimensional versions of) homogeneous Padé approximants applied to the Stieltjes transform.
- cubature formula at each point of grid
- solving a linear system of equations to retrieve the indicator function of K

・ 回 ト ・ ヨ ト ・ ヨ ト

#### Exact recovery.

- $\mathbf{G} = \{ x \in \mathbb{R}^n : g(\mathbf{x}) \le 1 \}$  compact.
- g is a nonnegative homogeneous polynomial
- Data are finitely many moments:

$$\mathbf{y}_{lpha} = \int_{\mathbf{G}} \mathbf{x}^{lpha} \, d\mathbf{x}, \quad lpha \in \mathbb{N}_{d}^{n}.$$

Also works for Quasi-homogeneous polynomials, i.e., when

$$g(\lambda^{u_1}x_1,\ldots,\lambda^{u_n}x_n) = \lambda g(x), \qquad x \in \mathbb{R}^n, \ \lambda > 0$$

for some vector  $\boldsymbol{u} \in \mathbb{Q}^n$ .

(*d*-Homogeneous =u-quasi homogeneous with  $u_i = \frac{1}{d}$  for all *i*).

ヘロン ヘアン ヘビン ヘビン

#### Exact recovery.

- $\mathbf{G} = \{ x \in \mathbb{R}^n : g(\mathbf{x}) \leq 1 \}$  compact.
- g is a nonnegative homogeneous polynomial
- Data are finitely many moments:

$$\mathbf{y}_{oldsymbol{lpha}}\,=\,\int_{\mathbf{G}}\mathbf{x}^{lpha}\,d\mathbf{x},\quad lpha\in\mathbb{N}_{d}^{n}.$$

Also works for Quasi-homogeneous polynomials, i.e., when

$$g(\lambda^{u_1}x_1,\ldots,\lambda^{u_n}x_n) = \lambda g(x), \qquad x \in \mathbb{R}^n, \, \lambda > 0$$

for some vector  $\boldsymbol{u} \in \mathbb{Q}^n$ .

(*d*-Homogeneous =u-quasi homogeneous with  $u_i = \frac{1}{d}$  for all *i*).

・ロト ・ 理 ト ・ ヨ ト ・

#### Exact recovery.

- $\mathbf{G} = \{ x \in \mathbb{R}^n : g(\mathbf{x}) \leq 1 \}$  compact.
- g is a nonnegative homogeneous polynomial
- Data are finitely many moments:

$$\mathbf{y}_{lpha}\,=\,\int_{\mathbf{G}}\mathbf{x}^{lpha}\,d\mathbf{x},\quad lpha\in\mathbb{N}_{d}^{n}.$$

Also works for Quasi-homogeneous polynomials, i.e., when

$$g(\lambda^{u_1}x_1,\ldots,\lambda^{u_n}x_n) = \lambda g(x), \qquad x \in \mathbb{R}^n, \, \lambda > 0$$

for some vector  $\boldsymbol{u} \in \mathbb{Q}^n$ .

(*d*-Homogeneous =u-quasi homogeneous with  $u_i = \frac{1}{d}$  for all *i*).

・ロ・ ・ 同・ ・ ヨ・ ・ ヨ・

#### Exact recovery.

- $\mathbf{G} = \{ x \in \mathbb{R}^n : g(\mathbf{x}) \leq 1 \}$  compact.
- g is a nonnegative homogeneous polynomial
- Data are finitely many moments:

$$\mathbf{y}_{lpha} = \int_{\mathbf{G}} \mathbf{X}^{lpha} \, d\mathbf{X}, \quad \mathbf{\alpha} \in \mathbb{N}_{d}^{n}.$$

Also works for Quasi-homogeneous polynomials, i.e., when

$$g(\lambda^{u_1}x_1,\ldots,\lambda^{u_n}x_n) = \lambda g(x), \qquad x \in \mathbb{R}^n, \, \lambda > 0$$

for some vector  $\boldsymbol{u} \in \mathbb{Q}^n$ .

(*d*-Homogeneous =u-quasi homogeneous with  $u_i = \frac{1}{d}$  for all *i*).

イロン 不良 とくほう 不良 とうほ

#### Exact recovery.

- $\mathbf{G} = \{ x \in \mathbb{R}^n : g(\mathbf{x}) \leq 1 \}$  compact.
- g is a nonnegative homogeneous polynomial
- Data are finitely many moments:

$$\mathbf{y}_{lpha} = \int_{\mathbf{G}} \mathbf{X}^{lpha} \, d\mathbf{X}, \quad \mathbf{\alpha} \in \mathbb{N}_{d}^{n}.$$

• Also works for Quasi-homogeneous polynomials, i.e., when

$$g(\lambda^{u_1}x_1,\ldots,\lambda^{u_n}x_n) = \lambda g(x), \qquad x \in \mathbb{R}^n, \ \lambda > 0$$

for some vector  $\boldsymbol{u} \in \mathbb{Q}^n$ .

(*d*-Homogeneous =u-quasi homogeneous with  $u_i = \frac{1}{d}$  for all *i*).

イロト イポト イヨト イヨト 三日

#### Exact recovery.

 G ⊂ ℝ<sup>n</sup> is open with G = int G and with real algebraic boundary ∂G. A polynomial of degree d vanishes on ∂G.

• Data are finitely many moments:

$$\mathbf{y}_{lpha} = \int_{\mathbf{G}} \mathbf{x}^{lpha} \, d\mathbf{x}, \quad lpha \in \mathbb{N}_{d}^{n}.$$

イロト イポト イヨト イヨト

э

- Exact recovery.
- G ⊂ ℝ<sup>n</sup> is open with G = int G and with real algebraic boundary ∂G. A polynomial of degree d vanishes on ∂G.
- Data are finitely many moments:

$$\mathbf{y}_{\alpha} = \int_{\mathbf{G}} \mathbf{x}^{\alpha} \, d\mathbf{x}, \quad \alpha \in \mathbb{N}_{d}^{n}.$$

・ 同 ト ・ ヨ ト ・ ヨ ト

ъ

- Exact recovery.
- G ⊂ ℝ<sup>n</sup> is open with G = int G and with real algebraic boundary ∂G. A polynomial of degree d vanishes on ∂G.
- Data are finitely many moments:

$$\mathbf{y}_{\alpha} = \int_{\mathbf{G}} \mathbf{x}^{\alpha} \, d\mathbf{x}, \quad \alpha \in \mathbb{N}_{d}^{n}.$$

・ 同 ト ・ ヨ ト ・ ヨ ト

ъ

So we are now concerned with PHFs, their sublevel sets and in particular, the integral

$$\mathbf{y} \mapsto \mathbf{I}_{g,h}(\mathbf{y}) := \int_{\{\mathbf{x} : \mathbf{g}(\mathbf{x}) \leq \mathbf{y}\}} h(\mathbf{x}) d\mathbf{x},$$

as a function  $I_{g,h} : \mathbb{R}_+ \to \mathbb{R}$  when g, h are PHFs.

#### With y fixed, we are also interested in

 $\boldsymbol{g}\mapsto \boldsymbol{I_{g,h}(y)},$ 

now as a function of g, especially when g is a nonnegative homogeneous polynomial.

Nonnegative homogeneous polynomials are particularly interesting as they can be used to approximate norms; see e.g. Barvinok

・ロト ・ 理 ト ・ ヨ ト ・

So we are now concerned with PHFs, their sublevel sets and in particular, the integral

$$\mathbf{y} \mapsto \mathbf{I}_{g,h}(\mathbf{y}) := \int_{\{\mathbf{x} : \mathbf{g}(\mathbf{x}) \leq \mathbf{y}\}} h(\mathbf{x}) d\mathbf{x},$$

as a function  $I_{g,h} : \mathbb{R}_+ \to \mathbb{R}$  when g, h are PHFs.

#### With y fixed, we are also interested in

 $\boldsymbol{g}\mapsto \boldsymbol{I_{g,h}(y)},$ 

now as a function of g, especially when g is a nonnegative homogeneous polynomial.

Nonnegative homogeneous polynomials are particularly interesting as they can be used to approximate norms; see e.g. Barvinok

イロト 不得 とくほ とくほ とう

As already observed in Morosov and Shakirov<sup>1</sup> the latter integral is related in a simple and remarkable manner to the non-Gaussian integral

 $\int_{\mathbb{R}^n} h \exp(-g) dx.$ 

Functional integrals appear frequently in quantum Physics

. ... where a challenging issue is to provide

exact formulas for  $\int \exp(-g) dx$ , the most well-known being when deg g = 2, i.e.,  $g(\mathbf{x}) = x^T Q x$ , with  $Q \succ 0$ ,

$$d = 2 \Rightarrow \int \exp(-g) \, dx = \frac{\operatorname{Cte}}{\sqrt{\operatorname{det}(Q)}}$$

Observe that det(Q) is an algebraic invariant of g,

<sup>1</sup>New and old results in Resultant theory, arXiv.0911.5278y1. I > I > OQ

As already observed in Morosov and Shakirov<sup>1</sup> the latter integral is related in a simple and remarkable manner to the non-Gaussian integral

$$\int_{\mathbb{R}^n} h \exp(-g) dx.$$

Functional integrals appear frequently in quantum Physics

.... where a challenging issue is to provide

exact formulas for  $\int \exp(-g) dx$ , the most well-known being when deg g = 2, i.e.,  $g(\mathbf{x}) = x^T Q x$ , with  $Q \succ 0$ ,

$$d = 2 \Rightarrow \int \exp(-g) \, dx = \frac{\operatorname{Cte}}{\sqrt{\operatorname{det}(Q)}}$$

Observe that det(Q) is an algebraic invariant of g,

<sup>1</sup>New and old results in Resultant theory, arXiv.0911.5278y1.. 🗉 📃 🧠 🦉

In particular, as a by-product in the important particular case when h = 1, they have proved that for all *forms g* of degree d,

$$\operatorname{Vol}\left(\{x : g(x) \le 1\}\right) = \int_{\{x : g(x) \le 1\}} dx$$
$$= \operatorname{cte}(d) \cdot \int_{\mathbb{R}^n} \exp(-g) d\mathbf{x},$$

where the constant depends only on *d* and *n*.

< 回 > < 三 >

In fact, a formula of exactly the same flavor was already known for convex sets, and was the initial motivation of our work. Namely, if  $C \subset \mathbb{R}^n$  is convex, its support function

$$x \mapsto \sigma_{\mathcal{C}}(x) := \sup \{x^T y : y \in \mathcal{C}\},\$$

is a PHF of degree 1, and the polar  $C^{\circ} \subset \mathbb{R}^n$  of *C* is the convex set  $\{x : \sigma_C(x) \leq 1\}$ .

Then ...

$$\operatorname{vol}(\mathcal{C}^\circ) = \frac{1}{n!} \int_{\mathbb{R}^n} \exp(-\sigma_{\mathcal{C}}(x)) \, dx, \qquad \forall \mathcal{C}.$$

▲ □ ▶ ▲ □ ▶ ▲

Let *g* be a nonnegative PHF such that  $vol({x : g(x) \le 1}) < \infty$ .

#### Theorem

Let g, h be PHFs of degree 0 < d and p respectively, then:

$$\int_{\{x:g(x)\leq y\}} h\,dx = \frac{y^{(n+p)/d}}{\Gamma(1+(n+p)/d)} \int_{\mathbb{R}^n} \exp(-g)\,h\,dx$$

$$\operatorname{vol}\left(\{x : g(x) \leq y\}\right) = \frac{y^{n/d}}{\Gamma(1+n/d)} \int_{\mathbb{R}^n} \exp(-g) \, dx$$

whenever the left-hand-side is finite.

◆□▶ ◆□▶ ◆三▶ ◆三▶ ● ● ●

### Proof

Observe that  $I_{g,h}(y)$  vanishes on  $(-\infty, 0]$ . For  $0 < \lambda \in \mathbb{R}$ , its Laplace transform  $\lambda \mapsto \mathcal{L}_{I_{g,h}}(\lambda) = \int_0^\infty \exp(-\lambda y) I_{g,h}(y) \, dy$  reads:

$$\mathcal{L}_{l_{g,h}}(\lambda) = \int_{0}^{\infty} \exp(-\lambda y) \left( \int_{\{x:g(x) \le y\}}^{\infty} h dx \right) dy$$
  
$$= \int_{\mathbb{R}^{n}} h(x) \left( \int_{g(x)}^{\infty} \exp(-\lambda y) dy \right) dx \quad \text{[by Fubini]}$$
  
$$= \frac{1}{\lambda} \int_{\mathbb{R}^{n}} h(x) \exp(-\lambda g(x)) dx$$
  
$$= \frac{1}{\lambda^{1+(n+p)/d}} \int_{\mathbb{R}^{n}} h(z) \exp(-g(z)) dz \quad \text{[by homog]}$$
  
$$= \frac{\int_{\mathbb{R}^{n}} h(z) \exp(-g(z)) dz}{\Gamma(1+(n+p)/d)} \mathcal{L}_{y^{(n+p)/d}}(\lambda).$$

# And so, by analyticity and the Identity theorem of analytical functions

$$I_{g,h}(\mathbf{y}) = \frac{\mathbf{y}^{(n+p)/d}}{\Gamma(1+(n+p)/d)} \int_{\mathbb{R}^n} h(x) \exp(-g(x)) dx,$$

< 🗇 🕨

프 🕨 🗉 프

- < ≣ → <

# I. Convexity

An interesting issue is to analyze how the Lebesgue volume  $\operatorname{vol} \{x \in \mathbb{R}^n : g(x) \leq 1\}$ , (i.e.  $\operatorname{vol} (G)$ ) changes with g.

#### Corollary

Let *h* be a PHF of degree *p* and let  $C_d \subset \mathbb{R}[x]_d$  be the convex cone of homogeneous polynomials *g* of degree at most *d* such that  $\int_G |h| dx < \infty$ . Then the function  $f_h : C_d \to \mathbb{R}$ ,

$$g\mapsto f_h(g):=\int_G h\,dx,\qquad g\in C_d,$$

- is a PHF of degree -(n + p)/d,
- convex whenever h is nonnegative and strictly convex if h > 0 on ℝ<sup>n</sup> \ {0}

イロト イポト イヨト イヨト

#### Corollary (continued)

Moreover, if h is continuous and  $g \in int(C_d)$  then:

$$\frac{\partial f_h(g)}{\partial g_\alpha} = \frac{-1}{\Gamma(1+(n+p)/d)} \int_{\mathbb{R}^n} x^\alpha h \exp(-g) dx$$
$$= \frac{-\Gamma(2+(n+p)/d)}{\Gamma(1+(n+p)/d)} \int_G x^\alpha h dx$$
$$\frac{\partial^2 f_h(g)}{\partial g_\alpha \partial g_\beta} = \frac{-1}{\Gamma(1+(n+p)/d)} \int_{\mathbb{R}^n} x^{\alpha+\beta} h \exp(-g) dx$$

イロト イポト イヨト イヨト 三日

#### **PROOF:** Just use

$$\int_{\{x: g(x) \le 1\}} h \, dx = \frac{1}{\Gamma(1 + (n + p)/d)} \int_{\mathbb{R}^n} h \exp(-g) \, dx$$

Notice that proving convexity directly would be non trivial but becomes easy when using the previous lemma!

Jean B. Lasserre Recovery of algebraic-exponential data from moments

・ 回 ト ・ ヨ ト ・ ヨ ト

э

#### **PROOF:** Just use

$$\int_{\{x: g(x) \le 1\}} h \, dx = \frac{1}{\Gamma(1 + (n + p)/d)} \int_{\mathbb{R}^n} h \exp(-g) \, dx$$

Notice that proving convexity directly would be non trivial but becomes easy when using the previous lemma!

For a set  $C \subset \mathbb{R}^n$ , recall:

• The support function  $x \mapsto \sigma_{C}(x) := \sup_{y \in C} \{x^{T}y : y \in C\}$ 

• The POLAR  $C^{\circ} := \{x \in \mathbb{R}^n : \sigma_C(x) \leq 1\}$ 

• and for a PHF *g* of degree *d*, its Legendre-Fenchel conjugate  $g^*(x) = \sup_{y} \{x^T y - g(y)\}$  is a PHF of degree *q* with  $\frac{1}{d} + \frac{1}{q} = 1$ .

個 とくほとくほとう ほ

#### Lemma

Let g be a closed proper convex PHF of degree 1 < d and let  $G = \{x : g(x) \le 1/d\}$ . Then:

$$G^{\circ} = \{x \in \mathbb{R}^{n} : g^{*}(x) \leq 1/q\}$$
  

$$vol(G) = \frac{p^{-n/p}}{\Gamma(1+n/p)} \int exp(-g) dx$$
  

$$vol(G^{\circ}) = \frac{q^{-n/q}}{\Gamma(1+n/q)} \int exp(-g^{*}) dx$$

 $\rightarrow$  yields completely symmetric formulas for g and its conjugate  $g^*$ .

イロト イポト イヨト イヨト 一臣

## Examples

• 
$$g(x) = |x|^3$$
 so that  $g^*(x) = \frac{2}{3\sqrt{3}}|x|^{3/2}$ . And so  
 $G = [-3^{-1/3}, 3^{-1/3}]; \quad G^\circ = [-3^{1/3}, 3^{1/3}].$ 

• TV screen: 
$$g(x) = x_1^4 + x_2^4$$
 so that  $g^*(x) = 4^{-4/3} \Im(x_1^{4/3} + x_2^{4/3})$ . And,

$$\mathbf{G} = \{x: x_1^2 + x_2^4 \leq \frac{1}{4}\}; \quad \mathbf{G}^\circ = \{x: x_1^{4/3} + x_2^{4/3} \leq 4^{1/3}\}.$$

• g(x) = |x| so that  $d \neq 1$ , and  $g^*(x) = 0$  if  $x \in [-1, 1]$ , and  $+\infty$  otherwise. Hence  $G = \{x : |x| \leq 1\} = [-1, 1]$  and with  $q = +\infty$ ,

$$G^{\circ} = [-1,1] = \{x : g^{*}(x) \leq \frac{1}{q} = 0\}.$$

◆□ > ◆□ > ◆臣 > ◆臣 > ─臣 ─のへで
If  $\mathbf{K} \subset \mathbb{R}^n$  is compact then computing the ellipsoid  $\xi$  of minimum volume containing **K** is a classical problem whose optimal solution is called the Löwner-John ellipsoid. So consider the following problem:

Find an homogeneous polynomial  $g \in \mathbb{R}[x]_{2d}$  such that its sub level set  $G := \{x : g(x) \le 1\}$  contains K and has minimum volume among all such levels sets with this inclusion property. Let  $\mathbf{P}[x]_{2d}$  be the convex cone of homogeneous polynomials of degree 2d whose sub-level set  $\mathbf{G} = \{x : g(x) \le 1\}$  has finite Lebesgue volume and with  $\mathbf{K} \subset \mathbb{R}^n$ , let  $C_{2d}(\mathbf{K})$  be the convex cone of polynomials nonnegative on  $\mathbf{K}$ .

#### Lemma

Let  $\mathbf{K} \subset \mathbb{R}^n$  be compact. The minimum volume of a sublevel set  $\mathbf{G} = \{\mathbf{x} : g(\mathbf{x}) \leq 1\}, g \in \mathbf{P}[x]_{2d}$ , that contains  $\mathbf{K} \subset \mathbb{R}^n$  is  $\rho/\Gamma(1 + n/2d)$  where:

$$\mathcal{P}: \qquad \rho = \inf_{g \in \mathbf{P}[x]_{2d}} \left\{ \int_{\mathbb{R}^n} \exp(-g) \, dx \, : \, 1 - g \, \in \, C_{2d}(\mathbf{K}) \right\}$$

a finite-dimensional convex optimization problem!

ヘロン 人間 とくほ とくほ とう

Let  $\mathbf{P}[x]_{2d}$  be the convex cone of homogeneous polynomials of degree 2d whose sub-level set  $\mathbf{G} = \{x : g(x) \le 1\}$  has finite Lebesgue volume and with  $\mathbf{K} \subset \mathbb{R}^n$ , let  $C_{2d}(\mathbf{K})$  be the convex cone of polynomials nonnegative on  $\mathbf{K}$ .

#### Lemma

Let  $\mathbf{K} \subset \mathbb{R}^n$  be compact. The minimum volume of a sublevel set  $\mathbf{G} = \{\mathbf{x} : g(\mathbf{x}) \leq 1\}, g \in \mathbf{P}[x]_{2d}$ , that contains  $\mathbf{K} \subset \mathbb{R}^n$  is  $\rho/\Gamma(1 + n/2d)$  where:

$$\mathcal{P}: \qquad \rho = \inf_{g \in \mathbf{P}[x]_{2d}} \left\{ \int_{\mathbb{R}^n} \exp(-g) \, dx \, : \, 1 - g \, \in \, C_{2d}(\mathbf{K}) \right\}$$

a finite-dimensional convex optimization problem!

イロト イ押ト イヨト イヨト

# Proof

• We have seen that:

$$\operatorname{vol}(\{x : g(x) \leq 1\}) = \frac{1}{\Gamma(1 + n/2d)} \int_{\mathbb{R}^n} \exp(-g) \, dx.$$

Moreover, the sub-level set  $\{x : g(x) \le 1\}$  contains **K** if and only if  $1 - g \in C_{2d}(\mathbf{K})$ , and so  $\rho/\Gamma(1 + n/2d)$  is the minimum value of all volumes of sub-levels sets  $\{x : g(x) \le 1\}$ ,  $g \in \mathbf{P}[\mathbf{x}]_{2d}$ , that contain **K**.

• Now since  $g \mapsto \int_{\mathbb{R}^n} \exp(-g) dx$  is strictly convex and  $C_{2d}(\mathsf{K})$  is a convex cone, problem  $\mathcal{P}$  is a finite-dimensional convex optimization problem.  $\Box$ 

< 回 > < 回 > < 回 > … 回

# III (continued). Characterizing an optimal solution

### Theorem

(a)  $\mathcal{P}$  has a unique optimal solution  $g^* \in \mathbf{P}[x]_{2d}$  and if  $g^* \in \operatorname{int}(\mathbf{P}[x]_{2d})$  there exists a Borel measure  $\mu^*$  supported on **K** such that:

(\*): 
$$\begin{cases} \int_{\mathbb{R}^n} x^{\alpha} \exp(-g^*) dx = \int_{\mathbf{K}} x^{\alpha} d\mu^*, \quad \forall |\alpha| = 2d \\ \int_{\mathbf{K}} (1 - g^*) d\mu^* = 0 \end{cases}$$

In particular,  $\mu^*$  is supported on the real variety  $V := \{x \in \mathbf{K} : g^*(\mathbf{x}) = 1\}$  and in fact,  $\mu^*$  can be substituted with another measure  $\nu^*$  supported on at most  $\binom{n+2d-1}{2d}$  points of V.

(b) Conversely, if  $g^* \in int(\mathbf{P}[x]_{2d})$  and  $\mu^*$  satisfy (\*) then  $g^*$  is an optimal solution of  $\mathcal{P}$ .

◆□▶ ◆□▶ ◆三▶ ◆三▶ ● ○ ○ ○

### Theorem

(a)  $\mathcal{P}$  has a unique optimal solution  $g^* \in \mathbf{P}[x]_{2d}$  and if  $g^* \in \operatorname{int}(\mathbf{P}[x]_{2d})$  there exists a Borel measure  $\mu^*$  supported on **K** such that:

(\*): 
$$\begin{cases} \int_{\mathbb{R}^n} x^{\alpha} \exp(-g^*) dx = \int_{\mathbf{K}} x^{\alpha} d\mu^*, \quad \forall |\alpha| = 2d \\ \int_{\mathbf{K}} (1 - g^*) d\mu^* = 0 \end{cases}$$

In particular,  $\mu^*$  is supported on the real variety  $V := \{x \in \mathbf{K} : g^*(\mathbf{x}) = 1\}$  and in fact,  $\mu^*$  can be substituted with another measure  $\nu^*$  supported on at most  $\binom{n+2d-1}{2d}$  points of V. (b) Conversely, if  $g^* \in \operatorname{int}(\mathbf{P}[x]_{2d})$  and  $\mu^*$  satisfy (\*) then  $g^*$  is an optimal solution of  $\mathcal{P}$ .

イロト イ理ト イヨト イヨト

3

Let  $\mathbf{K} \subset \mathbb{R}^2$  be the box  $[-1, 1]^2$ .

The set  $G_4 := \{x : g(x) \le 1\}$  with *g* homogeneous of degree 4 which contains **K** and has minimum volume is

$$\mathbf{x} \mapsto \mathbf{g}_4(\mathbf{x}) := x_1^4 + y_1^4 - x_1^2 x_2^2,$$

with  $vol(G_4) \approx 4.39$  much better than -  $\pi R^2 = 2\pi \approx 6.28$  for the Löwner-John ellipsoid of minimum volume, and

- the (convex) TV screen  $\textbf{G}:=\{\textbf{x}:(x_1^4+x_2^4)/2<=1\}$  with volume >5.

・ 同 ト ・ ヨ ト ・ ヨ ト



Jean B. Lasserre Recovery of algebraic-exponential data from moments

・ロト ・聞ト ・ヨト ・ヨト

E 900

Let  $\mathbf{K} \subset \mathbb{R}^2$  be the box  $[-1, 1]^2$ .

The set  $G_6 := \{x : g(x) \le 1\}$  with *g* homogeneous of degree 6 which contains **K** and has minimum volume is

$$\mathbf{x} \mapsto g_6(\mathbf{x}) := x_1^6 + y_1^6 - (x_1^4 x_2^2 + x_1^2 x_2^4)/2,$$

with  $vol(G_6) \approx 4.19$  much better than -  $\pi R^2 = 2\pi \approx 6.28$  for the Löwner-John ellipsoid of minimum volume, and

- better than the set  $G_4$  with volume 4.39.

▲帰▶ ▲ 国▶ ▲ 国▶



Jean B. Lasserre Recovery of algebraic-exponential data from moments

・ロト ・聞ト ・ヨト ・ヨト

₹ 990

# IV. Recovering g from moments of G

Write 
$$g(x) = \sum_{\beta} g_{\beta} x^{\beta}$$
.

#### Lemma

If g is nonnegative and d-homogeneous with G compact then:

$$\underbrace{\int_{G} x^{\alpha} g(x) dx}_{\sum_{\beta} g_{\beta} y_{\alpha+\beta}} = \frac{n+|\alpha|}{n+d+|\alpha|} \underbrace{\int_{G} x^{\alpha} dx}_{y_{\alpha}}, \qquad \alpha \in \mathbb{N}^{n}.$$

and so we see that the moments  $(y_{\alpha})$  satisfy linear relationships explicit in terms of the coefficients of the polynomial *g* that describes the boundary of *G*.

イロト イポト イヨト イヨト

So let us write  $\mathbf{g} \in \mathbb{R}^{s(d)}$  the unknown vector of coefficients of the unknown polynomial g.

Let  $\mathbf{M}_{d}(\mathbf{y})$  be the moment matrix of order d whose rows and columns are indexed in the canonical basis of monomials  $(x^{\alpha})$ ,  $\alpha \in \mathbb{N}_{d}^{n}$ , and with entries

 $\mathbf{M}_{\mathbf{d}}(\mathbf{y})(\alpha,\beta) = \mathbf{y}_{\alpha+\beta}, \qquad \alpha,\beta \in \mathbb{N}_{\mathbf{d}}^{n}.$ 

and let  $\mathbf{y}^d$  be the vector  $(\mathbf{y}_{\alpha}), \alpha \in \mathbb{N}^n_d$ .

Previous Lemma states that

 $\mathbf{M}_{\mathbf{d}}(\mathbf{y})\mathbf{g} = \mathbf{y}^{\mathbf{d}},$ 

or, equivalently,

$$\mathbf{g} = \mathbf{M}_{d}(\mathbf{y})^{-1} \, \mathbf{y}^{d},$$

because the moment matrix  $\mathbf{M}_d(\mathbf{y})$  is nonsingular whenever *G* has nonempty interior.

◆□ ▶ ◆□ ▶ ◆ □ ▶ ◆ □ ▶ ◆ □ ● ● ○ ○ ○

### In other words ...

one may recover g EXACTLY from knowledge of moments  $(y_{\alpha})$  of order d and 2d!

Jean B. Lasserre Recovery of algebraic-exponential data from moments

・ 4 回 ト ・ ( 回 ト ・

포 > 포

If g is not quasi-homogeneous then one cannot directly relate

$$\int_{\{\mathbf{x}:g(\mathbf{x})\leq 1\}} d\mathbf{x} \text{ and } \int_{\mathbb{R}^n} \exp(-g(\mathbf{x})) d\mathbf{x}.$$

But still the Laplace transform  $\lambda \mapsto F(\lambda)$  of the function

$$\mathbf{y} \mapsto f(\mathbf{y}) := \int_{\{\mathbf{x}: |g(\mathbf{x})| \le \mathbf{y}\}} d\mathbf{x}$$

is the non Gaussian integral

$$\lambda \mapsto F(\lambda) = \frac{1}{\lambda} \int_{\mathbb{R}^n} \exp(-\lambda |g(\mathbf{x})|) d\mathbf{x}.$$

🗇 🕨 🖉 🖻 🖌 🖉 🕨

Nice asymptotic results are available (Vassiliev)

$$f(\mathbf{y}) \approx \mathbf{y}^{\mathbf{a}} \ln(\mathbf{y})^{\mathbf{b}}, \text{ as } \mathbf{y} \rightarrow \infty$$

for some rationals *a*, *b* obtained from the Newton polytope of *g*.

One even has asymptotic results for

$$\mathbf{y} \mapsto \tilde{f}(\mathbf{y}) := \# \left( \{ \mathbf{x} : | \mathbf{g}(\mathbf{x}) | \le \mathbf{y} \} \cap \mathbf{Z}^n \right), \text{ as } \mathbf{y} \to \infty$$

## still in the form

$$ilde{f}({m y}) \,pprox\, {m y}^{a'}\,\ln({m y})^{b'}, \quad {
m as}\,\,{m y} \,{
m 
ightarrow} \,\infty$$

for some rationals a', b' obtained from the (modified) Newton polytope of g.

イロト イポト イヨト イヨト 一臣

Given a polynomial  $g \in \mathbb{R}[\mathbf{x}]_d$  write  $g(\mathbf{x}) = \sum_{k=0}^d g_k(\mathbf{x})$ , where each  $g_k$  is homogeneous of degree k.

### Lemma

Let  $g \in \mathbb{R}[\mathbf{x}]_d$  be such that its level set  $\mathbf{G} := {\mathbf{x} : g(\mathbf{x}) \le 1}$  is bounded. Then for every  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ :

$$\int_{\mathbf{G}} \mathbf{x}^{\alpha} (1 - g(\mathbf{x})) \, d\mathbf{x} = \sum_{k=1}^{d} \frac{k}{n + |\alpha|} \int_{\mathbf{G}} \mathbf{x}^{\alpha} g_{k}(\mathbf{x}) \, d\mathbf{x}$$

Observe that for each fixed arbitrary  $\alpha \in \mathbb{N}^n$  ..

One obtains LINEAR EQUALITIES between MOMENTS of the Lebesgue measure on G!

イロト イポト イヨト イヨト 三日

Given a polynomial  $g \in \mathbb{R}[\mathbf{x}]_d$  write  $g(\mathbf{x}) = \sum_{k=0}^d g_k(\mathbf{x})$ , where each  $g_k$  is homogeneous of degree k.

### Lemma

Let  $g \in \mathbb{R}[\mathbf{x}]_d$  be such that its level set  $\mathbf{G} := {\mathbf{x} : g(\mathbf{x}) \le 1}$  is bounded. Then for every  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ :

$$\int_{\mathbf{G}} \mathbf{x}^{\alpha} (1 - g(\mathbf{x})) \, d\mathbf{x} = \sum_{k=1}^{d} \frac{k}{n + |\alpha|} \int_{\mathbf{G}} \mathbf{x}^{\alpha} g_{k}(\mathbf{x}) \, d\mathbf{x}$$

Observe that for each fixed arbitrary  $\alpha \in \mathbb{N}^n \dots$ 

One obtains LINEAR EQUALITIES between MOMENTS of the Lebesgue measure on G!

イロト イポト イヨト イヨト 一日

# Proof:

Use Stokes' formula

$$\int_{\mathbf{G}} \operatorname{Div}(X) f(\mathbf{x}) \, d\mathbf{x} + \int_{\mathbf{G}} \langle X, \nabla f(\mathbf{x}) \rangle d\mathbf{x} = \int_{\partial \mathbf{G}} \langle X, \vec{n}_{\mathbf{x}} \rangle \, f \, d\sigma,$$

with vector field  $X = \mathbf{x}$  and  $f(\mathbf{x}) = \mathbf{x}^{\alpha}(1 - \mathbf{g}(\mathbf{x}))$ .

• Then observe that Div(X) = n and:

$$\langle X, \nabla f(\mathbf{x}) \rangle = |\alpha| f - \mathbf{x}^{\alpha} \sum_{k=1}^{d} k g_k(\mathbf{x}).$$

 $\star$  In the general case, when  $\partial G$  may have singular points, or lower dimensional components, we can invoke Sard's theorem, for the (smooth) sublevel sets

$$G_{\gamma} = \{ \mathbf{x} : g(\mathbf{x}) < \gamma \}$$

and pass to the limit  $\gamma o 1, \ \gamma < 1.$ 

# **Proof:**

• Use Stokes' formula

$$\int_{\mathbf{G}} \operatorname{Div}(X) f(\mathbf{x}) \, d\mathbf{x} + \int_{\mathbf{G}} \langle X, \nabla f(\mathbf{x}) \rangle d\mathbf{x} = \int_{\partial \mathbf{G}} \langle X, \vec{n}_{\mathbf{x}} \rangle \, f \, d\sigma,$$

with vector field  $X = \mathbf{x}$  and  $f(\mathbf{x}) = \mathbf{x}^{\alpha}(1 - g(\mathbf{x}))$ .

• Then observe that Div(X) = n and:

$$\langle X, \nabla f(\mathbf{x}) \rangle = |\alpha| f - \mathbf{x}^{\alpha} \sum_{k=1}^{d} k g_k(\mathbf{x}).$$

 $\star$  In the general case, when  $\partial G$  may have singular points, or lower dimensional components, we can invoke Sard's theorem, for the (smooth) sublevel sets

$$G_{\gamma} = \{\, \mathbf{x} : g(\mathbf{x}) < \gamma \,\}$$

and pass to the limit  $\gamma \rightarrow 1, \ \gamma < 1.$ 

イロン 不良 とくほう 不良 とうほ

Let  $\mathbf{G} \subset \mathbb{R}^n$  be open with  $\mathbf{G} = \operatorname{int} \overline{\mathbf{G}}$  and with real algebraic boundary  $\partial \mathbf{G}$ . A polynomial of degree *d* vanishes on  $\partial \mathbf{G}$ .

## Define a renormalised moment-type matrix $M_k^d(\mathbf{y})$ as follows:

- 
$$s(d) \ (= \binom{n+d}{n})$$
 columns indexed by  $\beta \in \mathbb{N}_d^n$ ,

- countably many rows indexed by  $\alpha \in \mathbb{N}_k^n$ , and with entries:

$$\mathbf{M}_{k}^{d}(\mathbf{y})(\alpha,\beta) := \frac{n+|\alpha|+|\beta|}{n+|\alpha|} \mathbf{y}_{\alpha+\beta}, \qquad \alpha \in \mathbb{N}_{k}^{n}, \, \beta \in \mathbb{N}_{d}^{n}.$$

▲ □ ▶ ▲ □ ▶ ▲

### Theorem

Let  $\mathbf{G} \subset \mathbb{R}^n$  be a bounded open set with real algebraic boundary. Assume that  $\mathbf{G} = \operatorname{int} \overline{\mathbf{G}}$  and a polynomial of degree d vanishes on  $\partial \mathbf{G}$  and not at 0. Then the linear system

$$\mathbf{M}_{2d}^{d}(\mathbf{y})\left[\begin{array}{c}-1\\\mathbf{g}\end{array}\right]=0,$$

admits a unique solution  $\mathbf{g} \in \mathbb{R}^{s(d)-1}$ , and the polynomial g with coefficients  $(0, \mathbf{g})$  satisfies

$$(\mathbf{x} \in \partial G) \Rightarrow (\underline{g}(\mathbf{x}) = 1).$$

(4個) (日) (日) (日) (日)

# Sketch of the proof

## The identity (obtained from Stokes' theorem)

$$\int_{\mathbf{G}} \mathbf{x}^{\alpha} (1 - g(\mathbf{x})) \, d\mathbf{x} = \sum_{k=1}^{d} \frac{k}{n + |\alpha|} \int_{\mathbf{G}} \mathbf{x}^{\alpha} g_{k}(\mathbf{x}) \, d\mathbf{x}$$

for all  $\alpha \in \mathbb{N}_k^n$ 

in fact reads:

$$\mathbf{M}_{k}^{d}(\mathbf{y})\left[\begin{array}{c}-\mathbf{1}\\\mathbf{g}\end{array}\right]=\mathbf{0},$$

Conversely, if g solves

$$\mathbf{M}_{2d}^{d}(\mathbf{y}) \left[ \begin{array}{c} -1 \\ \mathbf{g} \end{array} \right] = \mathbf{0},$$

then

$$\int_{\partial \mathbf{G}} \langle \mathbf{x}, \vec{n_{\mathbf{x}}} \rangle (1 - g(\mathbf{x})) \, \mathbf{x}^{\alpha} \, d\sigma = 0, \quad \forall \alpha \in \mathbb{N}^{n}_{2d}.$$

Jean B. Lasserre Recovery of algebraic-exponential data from moments

# Sketch of the proof

## The identity (obtained from Stokes' theorem)

$$\int_{\mathbf{G}} \mathbf{x}^{\alpha} (1 - g(\mathbf{x})) \, d\mathbf{x} = \sum_{k=1}^{d} \frac{k}{n + |\alpha|} \int_{\mathbf{G}} \mathbf{x}^{\alpha} g_{k}(\mathbf{x}) \, d\mathbf{x}$$

for all  $\alpha \in \mathbb{N}_k^n$ 

in fact reads:

$$\mathbf{M}_{k}^{d}(\mathbf{y})\left[\begin{array}{c}-1\\\mathbf{g}\end{array}\right]=\mathbf{0},$$

Conversely, if g solves

$$\mathbf{M}^{d}_{2d}(\mathbf{y}) \left[ \begin{array}{c} -1 \\ \mathbf{g} \end{array} \right] = \mathbf{0},$$

then

$$\int_{\partial \mathbf{G}} \langle \mathbf{x}, \vec{n_{\mathbf{x}}} \rangle (1 - g(\mathbf{x})) \, \mathbf{x}^{\alpha} \, d\sigma \, = \, \mathbf{0}, \quad \forall \alpha \in \mathbb{N}^{n}_{2d}.$$

Jean B. Lasserre

Recovery of algebraic-exponential data from moments

As  $\partial G$  is algebraic, one may write

$$ec{n_{\mathbf{x}}} = rac{
abla h(\mathbf{x})}{\|
abla h(\mathbf{x})\|},$$

for some polynomial h. Therefore

$$0 = \int_{\partial \mathbf{G}} \langle \mathbf{x}, \vec{n_{\mathbf{x}}} \rangle (1 - g(\mathbf{x})) \mathbf{x}^{\alpha} d\sigma \quad \forall \alpha \in \mathbb{N}_{2d}^{n}$$
  
$$= \int_{\partial \mathbf{G}} \underbrace{\langle \mathbf{x}, \nabla h(\mathbf{x}) \rangle}_{\in \mathbb{R}[\mathbf{x}]_{d}} \underbrace{(1 - g(\mathbf{x}))}_{\in \mathbb{R}[\mathbf{x}]_{d}} \mathbf{x}^{\alpha} \frac{1}{\|\nabla h\|} d\sigma \quad \forall \alpha \in \mathbb{N}_{2d}^{n}$$
  
$$\Rightarrow \int_{\partial \mathbf{G}} \underbrace{\langle \mathbf{x}, \nabla h(\mathbf{x}) \rangle^{2}}_{\neq 0 \ \sigma - a.e.} (1 - g(\mathbf{x}))^{2} d\sigma' = 0 \quad \Box$$

(過) (日) (

э

э

As  $\partial G$  is algebraic, one may write

$$ec{n_{\mathbf{x}}} = rac{
abla h(\mathbf{x})}{\|
abla h(\mathbf{x})\|},$$

for some polynomial h. Therefore

$$0 = \int_{\partial \mathbf{G}} \langle \mathbf{x}, \vec{n_{\mathbf{x}}} \rangle (1 - g(\mathbf{x})) \, \mathbf{x}^{\alpha} \, d\sigma \quad \forall \alpha \in \mathbb{N}_{2d}^{n}$$
  
$$= \int_{\partial \mathbf{G}} \underbrace{\langle \mathbf{x}, \nabla h(\mathbf{x}) \rangle}_{\in \mathbb{R}[\mathbf{x}]_{d}} \underbrace{(1 - g(\mathbf{x}))}_{\in \mathbb{R}[\mathbf{x}]_{d}} \, \mathbf{x}^{\alpha} \underbrace{\frac{1}{\|\nabla h\|} d\sigma}_{d\sigma'} \quad \forall \alpha \in \mathbb{N}_{2d}^{n}$$
  
$$\Rightarrow \int_{\partial \mathbf{G}} \underbrace{\langle \mathbf{x}, \nabla h(\mathbf{x}) \rangle^{2}}_{\neq 0 \, \sigma - a.e.} (1 - g(\mathbf{x}))^{2} d\sigma' = 0 \quad \Box$$

Jean B. Lasserre Recovery of algebraic-exponential data from moments

▲ □ ▶ ▲ □ ▶ ▲

프 🕨 🗉 프

For sake of rigor the boundary  $\partial G$  can be written

$$\partial \mathbf{G} = Z_0 \cup Z_1,$$

with  $Z_0$  being a finite union of smooth n - 1-submanifolds of  $\mathbb{R}^n$  leaving **G** on one side,  $Z_1$  is a union of the lower dimensional strata, and  $\sigma(Z_1) = 0$ .

・ 回 と ・ ヨ と ・ ヨ と

-

### Theorem

Let  $\mathbf{G} \subset \mathbb{R}^n$  be a bounded convex open set with real algebraic boundary. Assume that  $\mathbf{G} = \operatorname{int} \overline{\mathbf{G}}$ ,  $0 \in \mathbf{G}$ , and a polynomial of degree d vanishes on  $\partial \mathbf{G}$  and not at 0. Then the linear system

$$\mathbf{M}_d^d(\mathbf{y}) \left[ \begin{array}{c} -1 \\ g \end{array} \right] = \mathbf{0},$$

admits a unique solution  $\mathbf{g} \in \mathbb{R}^{s(d)-1}$ , and the polynomial g with coefficients  $(0, \mathbf{g})$  satisfies

$$(\mathbf{x} \in \partial G) \Rightarrow (\underline{g}(\mathbf{x}) = 1).$$

- 2

イロト イ理ト イヨト イヨト

 $\star$  As in the previous proof, if

$$\mathbf{M}_{d}^{d}(\mathbf{y})\left[egin{array}{c} -1\ g\end{array}
ight]=0,$$

then

$$\int_{\partial \mathbf{G}} \langle \mathbf{x}, \vec{n_{\mathbf{x}}} \rangle (1 - g(\mathbf{x}))^2 \, d\sigma \, = \, 0.$$

But one now uses that if  $0 \in \mathbf{G}$  then  $\langle \mathbf{x}, \vec{n_x} \rangle \ge 0$ .

・ 同 ト ・ ヨ ト ・ ヨ ト

э

## Example

Let us consider the two-dimensional example of the annulus

$$\textbf{G} \, := \, \{ \, \textbf{x} \in \mathbb{R}^2 : \, 1 - x_1^2 - x_2^2 \geq 0 ; \, x_1^2 + x_2^2 - 2/3 \geq 0 \, \}.$$



The polynomial  $(1 - x_1^2 - x_2^2)(x_1^2 + x_2^2 - 2/3)$  is the unique solution of  $\mathbf{M}_4^4(\mathbf{y})[-1, \mathbf{g}] = 0$ .

э

## Example continued: Non-algebraic boundary

Let 
$$\mathbf{G} = \{\mathbf{x} \in \mathbb{R}^2 : x_1 \ge -1; x_2 \ge 1; x_2 \le \exp(-x_1)\}.$$



≣ ) ≣

We now look as the eigenvector g of the smallest eigenvalue of  $M_3^3(y)$ .



Figure: Shape  $\mathbf{G}' = {\mathbf{x} : g(\mathbf{x}) \le 0}$  with d = 3

ъ

ъ

We now look as the eigenvector g of the smallest eigenvalue of  $\mathbf{M}_4^4(\mathbf{y})$ .



Figure: Shape  $\mathbf{G}' = {\mathbf{x} : g(\mathbf{x}) \le 0}$  with d = 4

ъ

ъ

uniformly supported on a set *G* of the form  $\{\mathbf{x} : g(\mathbf{x}) \le 1\}$ , for some polynomial  $g \in \mathbb{R}[\mathbf{x}]_d$ .

### Then :

• ALL moments  $y_{\alpha} := \int_{G} \mathbf{x}^{\alpha} d\mu$ ,  $\alpha \in \mathbb{N}^{n}$ , are determined from those up to order 3*d* (and 2*d* if *G* is convex) !

• A similar result holds true if now  $\mu$  has a density  $\exp(h(\mathbf{x}))$  on *G* (for some  $h \in \mathbb{R}[\mathbf{x}]$ ).

ightarrow is an extension to such measures of a well-known result for exponential families

(4回) (日) (日)

uniformly supported on a set *G* of the form  $\{\mathbf{x} : g(\mathbf{x}) \le 1\}$ , for some polynomial  $g \in \mathbb{R}[\mathbf{x}]_d$ .

## Then :

• ALL moments  $y_{\alpha} := \int_{G} \mathbf{x}^{\alpha} d\mu$ ,  $\alpha \in \mathbb{N}^{n}$ , are determined from those up to order 3*d* (and 2*d* if *G* is convex) !

• A similar result holds true

if now  $\mu$  has a density  $\exp(h(\mathbf{x}))$  on G (for some  $h \in \mathbb{R}[\mathbf{x}]$ ).

ightarrow is an extension to such measures of a well-known result for exponential families

ヘロト ヘワト ヘビト ヘビト

uniformly supported on a set *G* of the form  $\{\mathbf{x} : g(\mathbf{x}) \le 1\}$ , for some polynomial  $g \in \mathbb{R}[\mathbf{x}]_d$ .

## Then :

• ALL moments  $y_{\alpha} := \int_{G} \mathbf{x}^{\alpha} d\mu$ ,  $\alpha \in \mathbb{N}^{n}$ , are determined from those up to order 3*d* (and 2*d* if *G* is convex) !

• A similar result holds true if now  $\mu$  has a density  $\exp(h(\mathbf{x}))$  on *G* (for some  $h \in \mathbb{R}[\mathbf{x}]$ ).

ightarrow is an extension to such measures of a well-known result for exponential families

・ロ・ ・ 四・ ・ ヨ・ ・ ヨ・

uniformly supported on a set *G* of the form  $\{\mathbf{x} : g(\mathbf{x}) \le 1\}$ , for some polynomial  $g \in \mathbb{R}[\mathbf{x}]_d$ .

## Then :

• ALL moments  $y_{\alpha} := \int_{G} \mathbf{x}^{\alpha} d\mu$ ,  $\alpha \in \mathbb{N}^{n}$ , are determined from those up to order 3*d* (and 2*d* if *G* is convex) !

• A similar result holds true if now  $\mu$  has a density  $\exp(h(\mathbf{x}))$  on *G* (for some  $h \in \mathbb{R}[\mathbf{x}]$ ).

 $\rightarrow$  is an extension to such measures of a well-known result for exponential families

イロト イポト イヨト イヨト
- Compact sub-level sets  $G := \{x : g(x) \le y\}$  of homogeneous polynomials exhibit surprising properties. E.g.:
  - convexity of volume(G) with respect to the coefficients of g
  - Integrating a PHF *h* on *G* reduce to evaluating the non Gaussian integral  $\int h \exp(-g) dx$
  - A variational property yields a Gaussian-like property
  - exact recovery of *G* from finitely moments.
    (Also works for quasi-homogeneous polynomials with bounded sublevel sets!)
  - exact recovery for sets with algebraic boundary of known degree

・ロト ・ 同ト ・ ヨト ・ ヨト … ヨ

- COMPUTATION!: Efficient evaluation of  $\int_{\mathbb{R}^n} \exp(-g) dx$ , or equivalently, evaluation of vol  $(\{x : g(x) \le 1\}!$ 
  - The property

$$\int_{G} \mathbf{x}^{\alpha} g(x) \, dx = \frac{n + |\alpha|}{n + d + |\alpha|} \int_{G} x^{\alpha} \, dx, \qquad \forall \alpha,$$

helps a lot to improve efficiency of the method in Henrion, Lasserre and Savorgnan (SIAM Review)

▲ 同 ▶ ▲ 臣 ▶ ▲ 臣 ▶

- J.B. Lasserre. A Generalization of Löwner-John's ellipsoid Theorem. *Math. Program.*, to appear.
- J.B. Lasserre. Recovering an homogeneous polynomial from moments of its level set. *Discrete & Comput. Geom.* 50, pp. 673–678, 2013.
- J.B. Lasserre and M. Putinar. Reconstruction of algebraic-exponential data from moments. Submitted
- J.B. Lasserre. Unit balls of constant volume: which one has optimal representation? submitted.

## THANK YOU!

Jean B. Lasserre Recovery of algebraic-exponential data from moments

イロン イロン イヨン イヨン